

2 Sulfamed

Water Soluble Powder for Swine and Cattle

ACTIVE INGREDIENTS: Each gram contains: 630 mg of Sulfamethazine Sodium 315 mg of Sulfathiazole Sodium.

VETERINARY
USE ONLY

Net Contents
567 g/10 kg

DIN 02030586

2 Sulfamed is a strawberry flavoured, soluble mixture of the sodium salts of Sulfamethazine and Sulfathiazole for the treatment of bacterial diseases of cattle and swine that are susceptible to sulfonamides. A red dye is included in the product to alert users to the presence of sulfonamides.

Medprodex Inc.,
4-61 Malcolm Rd. Guelph, ON N1K 1A7



Indications:

Swine

As an aid in the treatment of enteritis and respiratory diseases caused by bacteria sensitive to sulfamethazine and sulfathiazole.

Cattle

As an aid in the treatment of bacterial infections such as pneumonia, metritis, enteritis, foot rot, mastitis and shipping fever complex.

Dosage & Administration:

Swine and Cattle

Herd medication: Dissolve 567 g in approximately 10 L of water. Add to each 340 L of drinking water the first day and to each 500 L on the second and subsequent days.

Individual Treatment for Cattle: Administer as a drench: On the first day, 60 mL of a 12.5% solution or 30 mL of a 25% solution per 35 kg of body weight, on the second, third and fourth day, administer half of the dosage of the first day.

How to Prepare Solution:

For a 12.5% solution: Dissolve 567 g in water to make a total volume of 4.5 L of solution.

For a 25% solution: Dissolve 567 g in water to make a total volume of 2.27 L of solution.
Use warm water and stir thoroughly.

NOTE

1 Tbsp. (15 mL) is equivalent to 7 g.

STORAGE

Store in a dry, cool place below 23°C.

CAUTION

If improvement is not seen within 48 hours, medication should be discontinued and diagnosis and the treatment re-evaluated. Generally, treatment should not exceed 4 days. Do not administer to animals with known sensitivity to sulfonamides or with renal impairments. Severely dehydrated animals should not be treated with this drug. Treated animals should have easy access to drinking water to maintain adequate hydration and miction.

WARNING(S)

Treated animals must not be slaughtered for use in food for at least 10 days after the latest treatment. Do not use in lactating dairy cattle. Do not administer in swine feed.

KEEP OUT OF REACH OF CHILDREN