

AMPROL[®] 25% FEED MIX

Veterinary
Use Only

Active Ingredient: amprolium 25% w/w

Net Contents 25 kg

DIN 00140503

[®]Trademark Huvepharma AD



RECOMMENDATIONS FOR USE IN TURKEYS

INDICATION: As an aid in the prevention of coccidiosis in turkeys.

USE DIRECTIONS: Use 0.0125% amprolium medicated feed as sole ration from one day of age up to and including sixteen (16) weeks of age.

RECOMMENDATIONS FOR USE IN BROILER CHICKENS

INDICATION: As an aid in the prevention of deaths from coccidiosis.

USE DIRECTIONS: Use 0.0125% or 0.025% amprolium medicated feed as sole ration from one day of age until birds are slaughtered.

RECOMMENDATIONS FOR USE IN REPLACEMENT CHICKENS

INDICATION (1): As an aid in the prevention of death from coccidiosis in chickens intended as conventional floor managed layers.

HEAVY EXPOSURE TO COCCIDIOSIS - Use 0.0125% - 0.008% amprolium medicated feed as sole ration from one day of age until 14 weeks as follows:

Age of chickens	Level of amprolium
0-5 weeks	0.0125%
5-9 weeks	0.0125% - 0.008%
9-14 weeks	0.008%

LIGHT TO MODERATE EXPOSURE TO COCCIDIOSIS - Use 0.008% - 0.006% amprolium medicated feed as sole ration from one day of age until 14 weeks as follows:

Age of chickens	Level of amprolium
0-5 weeks	0.008%
5-9 weeks	0.008% - 0.006%
9-14 weeks	0.006%

INDICATION (2):

As an aid in the prevention of deaths from coccidiosis in chickens intended as caged layers.

USE DIRECTIONS: Use 0.0125% or 0.025% amprolium medicated feed as sole ration from one day of age until birds are placed in cages.

RECOMMENDATIONS FOR USE IN CALVES

INDICATION: As an aid in the prevention and treatment of coccidiosis caused by Eimeria bovis and E. zuernii in calves.

USE DIRECTIONS:

PREVENTION - 250 mg of amprolium per 50 kg of body weight (5 mg/kg) per day for 21 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard.

TREATMENT - 500 mg of amprolium per 50 kg of body weight (10 mg/kg) per day for five (5) days.

The proper amount of amprolium medicated supplement is mixed with the amount of ration usually consumed in one day.

The table below gives suggested directions for using a 0.05% amprolium supplement.

	Animal weight (kg)	Supplement fed at rate of (kg/head/day)
For 21 day preventive program	100	1
	150	1.5
	200	2
	250	2.5
	300	3
For 5 day treatment program	100	2
	150	3
	200	4
	250	5
	300	6

RECOMMENDATIONS FOR MIXING: AMPROL should be thoroughly and evenly mixed in the feed in accordance with current good manufacturing practice for medicated feed. We recommend that a premix be made before the final mixing.

Expected level of amprolium in complete feed	Amount of AMPROL required per 1,000 kg of complete feed
0.0060%	240 g
0.0080%	320 g
0.0125%	500 g
0.0250%	1 kg
0.0500%	2 kg

WARNING:

CHICKENS AND TURKEYS: Do not feed to laying birds in production.

CALVES: Treated animals must not be slaughtered for use in food for at least seven (7) days after the latest treatment with this drug.

CAUTION:

CHICKENS AND TURKEYS:

1. Do not use amprolium medicated feed for treatment of outbreaks of coccidiosis.
2. If unexpected deaths occur, obtain an accurate diagnosis and follow veterinarian's or poultry pathologist's recommendations.

CALVES:

1. Do not use in feeds for calves intended for future breeding.
2. For a satisfactory diagnosis, a microscopic examination of the feces should be done by a veterinarian or diagnostic laboratory before treatment; when treating outbreaks, drug should be administered promptly after diagnosis is determined.
3. Do not use in feeds containing pellet binding agents except Pel-Aid.



AMPROL[®] 25% FEED MIX