

CHLOR 250 GRANULAR PREMIX

Veterinary Use Only

Net Weight 25 kg

DIN 00641979

Active Ingredients: chlortetracycline hydrochloride 44g/kg;
sulfamethazine 44g/kg; penicillin (from penicillin G procaine) 22g/kg



DIRECTIONS FOR USE

Pre starter and starter swine feeds:

Add and mix thoroughly into the ration at a rate to provide 2.5 kg CHLOR 250 per metric tonne (1000 kg) of finished (complete) feed. Chlortetracycline hydrochloride 0.011 % (110 mg/kg); sulfamethazine 0.011 % (110 mg/kg); penicillin 0.0055% (55 mg/kg).

1. As an aid in maintaining growth rate and feed efficiency in the presence of Atrophic Rhinitis - feed continuously as the sole ration: Note: This product has no effect on the organism causing Atrophic Rhinitis.
2. As an aid in maintaining weight gains and stimulating appetite during periods of stress caused by moving, vaccination, extreme temperature changes and castration - feed as sole ration during periods of stress and until 10 days after stress conditions have been eliminated.

Other Swine Feeds:

Note: This drug mixture is not to be used in grower or finisher rations for market hogs except on the basis of a veterinary prescription.

Add and mix thoroughly into the ration at a rate to provide 2.5 kg CHLOR 250 per metric tonne (1000 kg) of finished (complete) feed.

1. As an aid in maintaining growth rate and feed efficiency in the presence of Atrophic Rhinitis - feed continuously as the sole ration: Note: This product has no effect on the organism causing Atrophic Rhinitis.
2. As an aid in maintaining weight gains and stimulating appetite during periods of stress caused by moving, vaccination, extreme temperature changes and castration - feed as sole ration during periods of stress and until 10 days after stress conditions have been eliminated.
3. As an aid in the prevention of Bacterial Enteritis (including salmonellosis caused by *Salmonella choleraesuis*. and Vibrio Dysentery) - feed as sole ration during the period of early growth (25 to 30 kg).
4. As an aid in the treatment of Bacterial Enteritis (including salmonellosis caused by *Salmonella choleraesuis*. and Vibrio Dysentery) - feed continuously as the sole ration until three days after symptoms disappear.

Note: When treatment claim is made, labels must show this statement: Pigs off feed should be given individual treatment.

WARNING:

Treated animals must not be slaughtered for use in food for at least 10 days after the latest treatment with this drug.

STORAGE:

Store in a cool dry place below 30°C.
Keep from freezing