

CHLOR 50 GRANULAR PREMIX

Veterinary Use Only

Net Weight 25 kg

DIN 00641804

Active Ingredient: chlortetracycline hydrochloride 110g/kg



DIRECTIONS FOR USE

SWINE

0.011 % (110 mg/kg) of complete feed:

As an aid in maintaining weight gains and stimulating appetite during periods of stress caused by moving, vaccination, temperature changes and castration.

Feed as the sole ration during periods of stress and until 10 days after the stress condition has been eliminated.

As an aid in the treatment of bacterial enteritis (scours, bacterial diarrhea). Feed continuously as the sole ration following the appearance of symptoms and until 3 days after symptoms disappear.

Note:

1. Pigs off feed should be given individual treatment.
2. Pre-Starters and feeds to be fed up to and including 6 weeks of age are required to contain 0.022% chlortetracycline hydrochloride.

0.0055% (55 mg/kg) of complete feed:

As an aid in maintaining weight gains in the presence of Atrophic Rhinitis. Feed as the sole ration as long as symptoms of Atrophic Rhinitis persist.

Note:

1. This medicated feed has no effect on the organism causing Atrophic Rhinitis.
2. Not accepted for pre-starters.

As an aid in prevention of bacterial enteritis (scours, bacterial diarrhea). Feed as the sole ration during the period of early growth up to 32 kg (70lbs.)

Note:

Pre-Starters and feeds to be fed up to and including 6 weeks of age should contain 0.022% of chlortetracycline hydrochloride.

BEEF CATTLE

0.22 mg/kg of body weight/day or 70 mg/head/day in complete feed:

As an aid in the prevention of foot rot.

Feed continuously during the period animals are exposed to this disease.

CALVES UP TO 136 kg

0.0055% (55 mg/kg) of complete feed:

As an aid in the prevention of bacterial diarrhea in calves. Feed continuously during periods of early growth.

CALVES (Milk replacers)

0.0055% (55 mg/kg) of milk replacer powder.

As an aid in the prevention of bacterial diarrhea in calves. Feed continuously during periods of early growth.

WARNING:

Treated swine must not be slaughtered for use in food for at least 7 days after the latest treatment with this drug.

Treated cattle and calves must not be slaughtered for use in food for at least 5 days after the latest treatment with this drug.

Do not use in feed for dairy cows.

STORAGE:

Store in a cool dry place below 30°C.
Keep from freezing.



Bio Agri Mix LP
P.O. Box 399, Mitchell, ON
N0K 1N0
www.bioagrimix.com