

# COYDEN<sup>®</sup> 25

## ANTICOCCIDIAL MEDICATED PREMIX

Active Ingredient: Clopidol 250g/kg

Veterinary Use Only

Net Weight 25 kg

DIN 00656925

Huvepharma<sup>®</sup>, 33 James Boucher Blvd., Sofia 1407, Bulgaria



### Indications:

Coyden<sup>®</sup>25 coccidiostat is intended to be administered continuously in feed for the prevention of caecal and intestinal coccidiosis in broiler chickens and replacements for caged layers up to 16 weeks of age.

For use in the preparation of medicated feed for broiler chickens and replacements for caged layers

For feed manufacturing use only

### Dosage and Administration:

Broilers  
And  
Replacement  
Pullets

It is suggested that a mixture of COYDEN<sup>®</sup> 25 and some other feed ingredient be prepared before mixing in the finished ration. This will ensure thorough and even distribution of the feed. Use 500 grams of COYDEN<sup>®</sup> 25 per metric tonne of feed to produce feed containing 0.0125% clopidol. COYDEN<sup>®</sup> 25 should be blended thoroughly into the finished feed.

#### **FOR BROILER CHICKENS**

Feed containing clopidol should be fed continuously as the only ration from the time chicks are placed in floor pens until birds are slaughtered for meat.

#### **FOR REPLACEMENTS FOR CAGED LAYERS**

Feed containing clopidol should be fed continuously as the only ration up to 16 weeks of age.

#### **WARNING**

1. Do not feed to laying hens.
2. Do not feed clopidol to caged layer replacements after 16 weeks of age

#### **CAUTION**

1. Do not administer clopidol to birds intended for breeding purposes.
2. Consult a veterinarian, if losses exceed 0.5% in two days.
3. Do not use this medicated feed to suppress an outbreak of coccidiosis.

#### **NOTICE**

Do not ship or store with poisonous or high odour articles  
Do not contaminate

#### **STORAGE**

Store at room temperature in a dry place. Keep out of the reach of children.