

Oxytetracycline 100 GRANULAR PREMIX

Veterinary Use Only

Net Weight 25 kg

DIN 00654787

Active Ingredient: oxytetracycline hydrochloride 220g/kg



DIRECTIONS FOR USE

SPECIES	DISEASE OR CONDITION	INDICATIONS	kg of Oxy 100 PER TONNE COMPLETE FEED	FEEDING DIRECTIONS
SWINE	BACTERIAL ENTERITIS (scours)	As an aid in the prevention of bacterial enteritis.	1	Feed as the sole ration up to six weeks of age.
			0.25	Feed as the sole ration up to six weeks of age up to 27-32 kg body weight.
		As an aid in the treatment of bacterial enteritis.	1	Feed as the sole ration following the appearance of symptoms and until three days after symptoms disappear - for pigs up to six weeks of age.
			0.5	As above - for pigs greater than six weeks of age. NOTE: Pigs off feed should be given individual treatment.
	PRESENCE OF ATROPHIC RHINITIS	As an aid in maintaining weight gains in the presence of atrophic rhinitis. This product has no effect on the organism causing atrophic rhinitis.	0.25	Feed as the sole ration for as long as the symptoms of atrophic rhinitis persist. NOTE: Not acceptable for pre-starter feeds.
SWINE	STRESS	As an aid in stimulating appetite and maintaining weight gains during periods of stress caused by moving, vaccination, castration or extreme temperature changes.	0.5	Feed as the sole ration during periods of stress and until 10 days after the stress conditions have been eliminated.
	LEPTOSPIROSIS (abortion in swine)	As an aid in reducing the incidence of abortion caused by leptospirosis. NOTE: Good sanitation procedures must be practiced to prevent spread of infections.	2.5	When abortion in swine has been diagnosed as due to leptospirosis, feed at once as the sole ration to pregnant and affected sows for a two-week period.
BEEF CATTLE	BLOAT	As an aid in reducing the incidence of bloat in young cattle on pasture and in feed lots.	Feed continuously. Level of oxytetracycline hydrochloride in the complete feed is to be such that each animal will receive 75 mg per day. 55 g of oxytetracycline hydrochloride (0.25 kg of oxytetracycline 100) per tonne complete feed when fed at the rate of 1.36 kg per head per day meets the 75 mg level stated above.	
CALVES	BACTERIAL ENTERITIS (scours)	As an aid in the prevention of bacterial enteritis.	0.25	Feed continuously during the period of early growth (up to 136 kg body weight).
	STRESS	As an aid in stimulating appetite and maintaining weight gains or egg production, hatchability and feed efficiency during periods of stress caused by moving, debeaking, vaccination or extreme temperature changes.	0.5	Feed as the sole ration. In case of vaccination, feed for 8-10 days immediately following vaccination. For other stress conditions, feed for at least five days.
TURKEYS and/or CHICKENS (broilers, replacement birds or layers)	HISTORY OF C.R.D. and/or NON-SPECIFIC ENTERITIS ON PREMISES	As an aid in stimulating appetite and maintaining weight gains when chickens are expected to be exposed to mild attacks of C.R.D. and/or non-specific enteritis.	0.25	Feed as the sole ration for as long as the birds are expected to be exposed to the infection.
	PRESENCE OF C.R.D.	As an aid in stimulating appetite and maintaining weight gains or egg production, hatchability, eggshell quality and feed efficiency in the presence of C.R.D.	0.5	Feed as the sole ration from the onset of the disease for at least one month or until a few days after the symptoms disappear.
	PRESENCE OF C.R.D. and STRESS	As an aid in stimulating appetite and maintaining weight gains or egg production in the presence of C.R.D. and while under stress caused by moving, debeaking, vaccination or extreme temperature changes.	1	Feed as the sole ration from the onset of the disease for at least one month or until a few days after the symptoms disappear.
	PRESENCE OF NON-SPECIFIC ENTERITIS	As an aid in stimulating appetite and maintaining weight gains in the presence of non-specific enteritis.	0.5	Feed as the sole ration from the onset of the disease for at least one month or until a few days after the symptoms disappear.
	INFECTIOUS SINUSITIS	As an aid in the prevention of infectious sinusitis of turkeys.	0.5	Feed as the sole ration for as long as the birds are expected to be exposed to the infection.
TURKEYS and/or CHICKENS (broilers, replacement birds or layers)	INFECTIOUS SYNOVITIS	As an aid in the prevention of infectious synovitis of turkeys.	0.5	Feed as the sole ration for as long as the birds are expected to be exposed to the infection.
		As an aid in the treatment of infectious synovitis of turkeys.	1	Feed as the sole ration when symptoms appear and continue until at least two weeks after symptoms disappear.
LAMBS	BACTERIAL ENTERITIS (scours)	As an aid in the reduction of bacterial enteritis in creep-fed suckling lambs.	0.5	Feed as the sole creep feed during the suckling period.
	ENTEROTOXEMIA (overeating disease)	As an aid in the reduction of losses due to enterotoxemia in feedlot lambs.	0.1	Feed as the sole ration while lambs are in the feedlot

WARNING Treated animals must not be slaughtered for use in food for at least 7 days for swine, chickens and turkeys, 5 days for cattle and calves, and 4 days for lambs after the latest treatment with this drug. Do not administer to lactating dairy cattle.

STORAGE Store in a cool dry place below 30°C. Keep from freezing.



Bio Agri Mix LP
P.O. Box 399, Mitchell, ON
N0K 1N0
www.bioagrimix.com