

Oxy 250

Soluble Antibiotic for Chickens, Turkeys, Swine, Calves, Sheep and Lambs

ACTIVE INGREDIENTS: Each gram contains 250 mg Oxytetracycline HCl USP.


VETERINARY
USE ONLY

Package sizes
100 g/10 kg

DIN 00780782

Oxytetracycline HCl is an antibiotic similar to tetracycline and chlortetracycline in bacterial spectrum. Oxytetracycline is rapidly absorbed and effective in the treatment of many diseases due to gram-positive and gram-negative bacteria, spirochetes, rickettsia, and some chlamydia.

Medprodex Inc.,
4-61 Malcolm Rd. Guelph, ON N1K 1A7



Indications:

As an aid in the treatment of infections caused by or associated with micro-organisms sensitive to oxytetracycline HCl.

Chickens and Turkeys

C.R.D., bluecomb, cholera, enteritis and hexamitiasis.

Swine, Calves, Sheep
and Lambs

Enteritis (diarrhea) and respiratory diseases.

Dosage & Administration:

Chickens and Turkeys

1 g of powder per 5 L of drinking water or 100 g per 500 L of water for 3 to 5 days.

For automatic proportioner: Set the apparatus to distribute 30 mL per 4 L of drinking water (1 oz/ gallon U.S.). Prepare the stock solution by dissolving 100 g in 3.8 L of water. This will medicate 500 L of drinking water.

Swine

1 g of powder per 25 kg of body weight every 12 hours, given as a drench using a dose syringe or 2 g of powder per 4 L of drinking water or 100 g per 200 L of drinking water for 4 or 5 days.

For automatic proportioner: Set the apparatus to distribute 30 mL per 4 L of drinking water (1 oz/ gallon U.S.). Prepare the stock solution by dissolving 200 g in 3 L of water. This will medicate 400 L of drinking water.

Calves (scours)

1 g of powder per 50 kg of body weight every 12 hours, given as a drench in water or 100 g per 225 L of water for 3 to 5 days.

Calves (respiratory diseases)

Double the above dose.

Sheep and Lambs

1 g of powder per 25 kg of body weight every 12 hours, given as a drench in water for 4 to 5 days.

NOTE

Only medicated drinking water should be provided during the treatment period and fresh solutions should be prepared daily. 1 teaspoon (5 mL) equals 4 g.

STORAGE

Store in a dry, cool place below 23°C.

WARNING(S)

Treated animals and birds must not be slaughtered for use in food for at least 7 days for chickens and turkeys; 10 days for swine, calves and lambs after the latest treatment with this drug. Do not feed to laying birds. Do not use in lactating dairy animals.

KEEP OUT OF REACH OF CHILDREN