

SACOX[®] 120

ANTICOCCIDIAL MEDICATED PREMIX

Active Ingredient :Each kg contains 120 g of salinomycin

Veterinary Use Only

Net Weight 25 kg

DIN 02230508

Huvepharma[®], 33 James Boucher Blvd., Sofia 1407, Bulgaria



Indications:

SACOX[®] 120 premix is indicated as an aid in the prevention of coccidiosis caused by *Eimeria acervulina*, *E. mitis*, *E. necatrix*, *E. maxima*, *E. tenella* and *E. brunetti* in broiler chickens.

Broilers

For use in complete feeds only
For feed manufacturing use only

Dosage and Administration:

Thoroughly mix 500 grams (0.5 kg) of SACOX[®] 120 Premix with 999.5 kg of complete feed to obtain a finished feed containing 60 mg / kg (60 ppm) salinomycin sodium. The medicated feed is to be fed continuously as the sole ration up until slaughter.

Feed this medicated feed as the sole ration.

WARNING

1. Do not feed to replacement, laying or breeding chickens.
2. Operators handling the concentrated product should wear a face mask, gloves and protective clothing.
3. Keep this, and all drugs, out of the reach of children.

CAUTION

1. Do not allow turkeys, dogs or horses access to this medicated feed as it is known to be toxic to these species. Extra care should be taken to avoid contamination of feeds for these animals.
2. A weight gain reduction may occur in cockerels fed this medicated feed in the absence of coccidial exposure.
3. Do not administer tiamulin to birds receiving this medicated feed.
4. Do not use in feeds containing pellet binding agents
5. Do not use this feed for treatment of outbreaks of coccidiosis.
6. Consult a veterinarian or poultry pathologist if losses exceed 0.5% in a two day period.

STORAGE

Store at room temperature in a dry place. Keep out of the reach of children.