

# SALINOMYCIN 60

## ANTICOCCIDIAL MEDICATED PREMIX

Veterinary Use Only

Net Weight 25 kg

DIN 02237431

Active Ingredient: salinomycin sodium 60g/kg



### Indications:

Salinomycin 60 Premix is indicated as an aid in the prevention of coccidiosis caused by *Eimeria acervulina*, *E. mitis*, *E. necatrix*, *E. maxima*, *E. tenella*, and *E. brunetti*, in broiler chickens.

Broilers:

For feed manufacturing use only

### Directions for use:

Thoroughly mix 1000 grams (1.0 kg) of Salinomycin 60 Premix with 999 kg of complete feed to obtain a finished feed containing 60 mg/kg (60 ppm) salinomycin sodium. The medicated feed is to be fed continuously as the sole ration up until slaughter.

### WARNING:

1. Do not feed to replacement, laying or breeding chickens.
2. Operators handling the concentrated product should wear a face mask, gloves and protective clothing.
3. Keep this and all drugs out of the reach of children.

### CAUTION:

1. Do not allow turkeys, dogs or horses access to this medicated feed as it is known to be toxic to these species. Extra care should be taken to avoid contamination of feeds for these animals.
2. A weight gain reduction may occur in cockerels fed this medicated feed in the absence of coccidial exposure.
3. Do not administer tiamulin to birds receiving this medicated feed.
4. Do not use in feeds containing pellet binding agents.
5. Do not use this feed for treatment of outbreaks of coccidiosis.
6. Consult a veterinarian or poultry pathologist if losses exceed 0.5% in a two day period.
7. Do not leave or dispose of product where it may be accessible to animals not under treatment and for wildlife.

### STORAGE:

Store in a cool, dry place below 30°C.  
Keep from freezing.



Bio Agri Mix LP  
P.O. Box 399, Mitchell, ON  
N0K 1N0  
[www.bioagrimix.com](http://www.bioagrimix.com)