

Tetramed 1000

Soluble Antibiotic for **Chickens, Turkeys, Swine, Calves, Sheep and Lambs**

ACTIVE INGREDIENTS: Each gram contains 1000 mg Tetracycline HCl USP.

VETERINARY
USE ONLY

Package sizes
500 g/2.5 kg/10 kg

DIN 00894486

Medprodex Inc.,
4-61 Malcolm Rd. Guelph, ON N1K 1A7



Indications:

As an aid in the treatment of infections caused by or associated with micro-organisms sensitive to tetracycline HCl.

Chickens and Turkeys

C.R.D., bluecomb, cholera, enteritis and hexamitiasis.

Swine, Calves, Sheep
and Lambs

Enteritis (diarrhea) and respiratory diseases.

Dosage & Administration:

Chickens and Turkeys

1 g of powder per 20 L of drinking water or 100 g per 2000 L of water for 3 to 5 days.

For automatic proportioner: Set the apparatus to distribute 30 mL per 4 L of drinking water (1 oz/gallon U.S.). Prepare the stock solution by dissolving 25 g of powder in 4 L of water. This will medicate 500 L of drinking water.

Swine (scours and
respiratory diseases)

1 g of powder per 100 kg of body weight every 12 hours, given as a drench using a dose syringe or 1 g of powder per 8 L of drinking water or 25 g of powder per 200 L of drinking water for 4 or 5 days.

For automatic proportioner: Set the apparatus to distribute 30 mL per 4 L of drinking water. Prepare the stock solution by dissolving 50 g of powder in 3 L of water. This will medicate 400 L of drinking water.

Calves (scours)

0.25 g of powder per 50 kg of body weight every 12 hours, given as a drench in water or 25 g of powder per 225 L of water for 3 to 5 days.

Calves (respiratory diseases)

Double the above dose.

Sheep and Lambs
(scours and respiratory diseases)

0.25 g of powder per 25 kg of body weight every 12 hours, given as a drench in water for 4 or 5 days.

NOTE

Only medicated drinking water should be provided during the treatment period and fresh solutions should be prepared daily. 1 teaspoonful (5 mL) equals 4 g. 1 scoop (43 mL) equals 25 g.

STORAGE

Store in a dry, cool place below 23°C.

WARNING(S)

Treated animals and birds must not be slaughtered for use in food for at least 5 days after the latest treatment with this drug. Do not feed to laying birds. Do not use in lactating dairy cattle.

KEEP OUT OF REACH OF CHILDREN