

Tetramed 250

Soluble Antibiotic for **Chickens, Turkeys, Swine, Calves, Sheep and Lambs**

ACTIVE INGREDIENTS: Each gram contains 250 mg Tetracycline HCl USP.

VETERINARY
USE ONLY

Package sizes
400 g/10 kg

DIN 00780863

Tetracycline HCl is an antibiotic similar to oxytetracycline and chlortetracycline in bacterial spectrum. Tetracycline is rapidly absorbed and effective in the treatment of many diseases due to gram-positive and gram-negative bacteria, spirochetes, rickettsia, and some chlamydia.

Medprodex Inc.,
4-61 Malcolm Rd. Guelph, ON N1K 1A7



Indications:

As an aid in the treatment of infections caused by or associated with micro-organisms sensitive to tetracycline HCl.

Chickens and Turkeys

As an aid in the treatment of C.R.D., bluecomb, cholera, enteritis and hexamitiasis.

Swine, Calves, Sheep
and Lambs

As an aid in the treatment of enteritis (diarrhea) and respiratory diseases.

Dosage & Administration:

Chickens and Turkeys

1 g of powder per 5 L of drinking water or 100 g per 500 L or 400 g per 2000 L of water for 3 to 5 days.

For automatic proportioner: Set the apparatus to distribute 30 mL per 4 L of drinking water (1 oz/gallon U.S.). Prepare the stock solution by dissolving 100 g in 3.8 L of water. This will medicate 500 L of drinking water or set the apparatus to distribute 15 mL per 4 L of water (1/2 oz/gallon U.S.). Prepare the stock solution by dissolving 400 g in 7.5 L of water. This will medicate 2,000 L of drinking water.

Swine

1 g of powder per 25 kg of body weight every 12 hours, given as a drench using a dose syringe or 2 g of powder per 4 L of drinking water or 100 g per 200 L of drinking water or 400 g per 800 L for 4 or 5 days.

For automatic proportioner: Set the apparatus to distribute 30 mL per 4 L of drinking water (1 oz/gallon U.S.). Prepare the stock solution by dissolving 200 g in 3 L of water or 400 g in 6 L of water. This will medicate 400 L and 800 L of drinking water.

Calves (scours)

1 g of powder per 50 kg of body weight every 12 hours, given as a drench in water or 100 g of powder per 225 L of water or 400 g per 900 L of water for 3 to 5 days.

Calves (respiratory diseases)

Double the above dosage.

Sheep and Lambs

1 g of powder per 25 kg of body weight every 12 hours, given as a drench in water for 4 or 5 days.

NOTE

Only medicated drinking water should be provided during the treatment period and fresh solutions should be prepared daily. 1 teaspoon (5 mL) equals 4 g.

STORAGE

Store in a dry, cool place below 23°C.

WARNING(S)

Treated animals and birds must not be slaughtered for use in food for at least 5 days after the latest treatment with this drug. Do not feed to laying birds. Do not use in lactating dairy cattle.

KEEP OUT OF REACH OF CHILDREN