

# TIAMULIN 1.78% PREMIX

Veterinary Use Only

Net Weight 25 kg

DIN 02242446

Active Ingredient: tiamulin base 17.8g/kg  
(equivalent to 22g tiamulin hydrogen fumarate per kg)



**Indications** For use in swine for the treatment and for the prevention of swine dysentery associated with *Serpulina (Treponema) hyodysenteriae* susceptible to tiamulin.

**Directions** Thoroughly mix TIAMULIN 1.78% PREMIX into complete nonmedicated swine feed for the treatment or for the prevention of swine dysentery at the rates per metric tonne of feed as shown in the table below.

Indication	Tiamulin base use level	Amount of TIAMULIN 1.78% PREMIX per tonne of feed
Treatment	178.1 g/tonne	10.0 kg
Prevention	31.2 g/tonne	1.75 kg

**Treatment** - Feed continuously as the sole ration at 178.1 g tiamulin base/tonne for 14 days.

**Prevention** - Feed continuously at 31.2 g tiamulin base/tonne as the sole ration to swine on premises with a history of swine dysentery but where signs of disease have not yet occurred. Feed continuously at 31.2 g tiamulin base/tonne to prevent reinfection following use of tiamulin at 178.1 g tiamulin base/tonne for 14 days or following use of tiamulin medicated drinking water for treatment of swine dysentery associated with *Serpulina (Treponema) hyodysenteriae*.

**WARNING**

1. Treated animals must not be slaughtered for use in food for at least 7 days after the latest treatment with this drug at 178.1 g tiamulin base/tonne for treatment of swine dysentery or at least 2 days after use at 31.2g tiamulin base/tonne for prevention of swine dysentery.
2. Direct contact with skin or mucous membranes may cause irritation. If contact occurs, wash with soap and water.

**CAUTION**

1. Do not feed to animals other than swine.
2. Do not feed undiluted.
3. Swine being treated with tiamulin should not have access to or be treated with tiamulin incompatible polyether ionophores (e.g.: monensin, salinomycin, narasin).
4. Use as the only source of tiamulin. Discontinue use if signs of toxicity occur.
5. Do not use in feeds containing pellet binding agents.
6. Do not feed to gilts and sows during 4 weeks after service.

**ADVERSE REACTIONS** (Associated with the use of 178.1 g tiamulin base/tonne for treatment of swine dysentery).

1. Overdoses of tiamulin have sometimes produced transitory salivation, vomiting and an apparent calming effect on the pig.
2. In rare cases, redness of the skin, primarily over the ham and underline, has been observed during medication.
3. If signs of toxicity or redness of the skin occurs, promptly discontinue the use of the therapeutic level of medicated feed.

Note: During treatment, swine should be housed under conditions of adequate space and sanitation.

**STORAGE** Store in a cool dry place below 30°C.  
Keep from freezing



Bio Agri Mix LP  
P.O. Box 399, Mitchell, ON  
N0K 1N0  
www.bioagrimix.com