

VIRGINIAMYCIN 44


MEDICATED PREMIX

Veterinary Use Only

Net Weight 25 kg

DIN 02215691

Active Ingredient: virginiamycin 44g/kg



Broiler
Chickens
and swine

Indications:

For increased weight of gain in swine and broiler chickens.
For improved feed efficiency in broiler chickens.
As an aid in the treatment and control of swine dysentery.
As aid in the prevention of necrotic enteritis in broiler chickens caused by *Clostridium perfringens* susceptible to virginiamycin.

Use directions:

VIRGINIAMYCIN 44 PREMIX must be thoroughly mixed in the feed to assure even distribution. When the required amount is less than 5 kg per 1000 kg of complete feed, dilute it with one of the feed ingredients (enough to make a quantity equal to 10 kg) before final mixing.

DOSAGE

REQUIRED LEVEL OF VIRGINIAMYCIN ACTIVITY PER 1000 KG OF COMPLETE FEED	REQUIRED AMOUNT OF VIRGINIAMYCIN 44 PREMIX PER 1000 KG OF COMPLETE FEED	APPROXIMATE AMOUNT OF VIRGINIAMYCIN ACTIVITY PROVIDED PER KG BODY WEIGHT PER ANIMAL PER DAY
SWINE		
For treatment and control of swine dysentery:		
110 grams for two weeks followed by:	2.5 kg	5.00 mg
55 grams for 4 weeks	1.25 kg	2.50 mg
After cessation of medication at 55 grams per 1000 kg, if swine dysentery recurs, re-treat with the above regimen		
For increased rate of weight gain*:		
11 grams	0.25kg	0.50mg
* START MEDICATION AT WEANING AND CONTINUE TO FEED THROUGHOUT GROWING AND FINISHING PHASES UNTIL SWINE REACH MARKET WEIGHT.		
BROILER CHICKENS		
For increased rate of weight gain and improved feed efficiency:		
11 grams	0.25 kg	0.50 mg
For the prevention of necrotic enteritis caused by <i>Clostridium perfringens</i> susceptible to virginiamycin:		
22 grams	0.50 kg	1.00 mg
FEED ONLY TO BROILER CHICKENS FROM DAY ONE TO MARKET WEIGHT		

WARNING

Do not feed to birds producing eggs for human consumption.

CAUTION

1. Do not feed to replacement or breeding chickens
2. Compatible with bentonite, Pel-Aid, Agri-Colloid, or Lignosol, when used as pellet binding agents.
3. Store in a cool dry place below 30°C. Keep from freezing.
4. Close container after each use.



Bio Agri Mix LP
P.O. Box 399, Mitchell, ON
N0K 1N0
www.bioagrimix.com