

ZOAMIX[®]

ANTICOCCIDIAL MEDICATED PREMIX

Active Ingredient: zoalene 250g/kg

Veterinary Use Only

Net Weight 25 kg

DIN 00656917

Huvepharma[®], 33 James Boucher Blvd., Sofia 1407, Bulgaria



Indications:

As an aid in the prevention of caecal and intestinal coccidiosis in broiler chickens, replacement chickens and growing turkeys.

For use in complete feeds only

Use directions:

It is suggested that a mixture of ZOAMIX and some other feed ingredient be prepared prior to mixing in with the finished ration. This will ensure thorough and even distribution of the drug in the medicated feed.

For Broiler Chickens: Use 500 grams of ZOAMIX per metric tonne of finished product to produce a medicated feed containing 0.0125% w/w zoalene. ZOAMIX should be thoroughly blended into the finished medicated feed. Feed medicated with zoalene should be fed continuously as the sole ration from the time chicks are placed in floor pens until they are slaughtered for meat.

For Replacement Chickens: Feed medicated with zoalene, when used under conditions of exposure to coccidiosis, will allow immunity to develop to protect against losses due to the disease when the birds are placed on non-medicated feed for egg laying purposes. Use 500 grams of ZOAMIX per metric tonne of finished product to produce a medicated feed containing 0.0125% w/w of zoalene. Feed medicated with zoalene should be fed continuously as the sole ration from day one. Discontinue feed medicated with zoalene when birds are 14 weeks old.

Turkeys: When turkey poults are reared in confinement and coccidiosis is usually a problem, use 750 grams ZOAMIX per metric tonne of finished feed to produce a medicated feed containing 0.0187% w/w zoalene. Feed medicated with zoalene should be fed continuously until the birds are 8 weeks of age, and when local conditions warrant, may be fed up to 14 weeks of age.

Chickens
and
turkeys

DIRECTIONS

Feed this medicated feed as the sole ration.

WARNING

1. Do not feed to laying chickens.
2. Do not feed to laying turkey hens.
3. Do not feed to replacement chickens after the first 14 weeks of life

STORAGE

Store at room temperature in a dry place. Keep out of the reach of children.

CAUTION

1. Do not feed this medicated feed for the treatment of outbreaks of coccidiosis.
2. If unexpected deaths occur consult a veterinarian or poultry pathologist and follow his instructions