Pot-Pen® 15 000 000 000 I.U.

INGREDIENT(S):

penicillin G potassium

USE:

Necrotic enteritis (Clostridium perfringens), control/prevention

SPECIES:

Chickens

ROA:

dw

DOSE:

1 jar/1,682 L (370 Imperial gal) drinking water (0.2 g/L drinking water).

1 pail/50,460 L (11,100 Imperial gal) drinking water (20 g/L drinking water).

(Provides ~297,000 I.U. penicillin/ L).

Give as sole source of drinking water for 5d.

WITHDRAWAL TIME:

Meat: 1d

USE:

Streptococcus suis infections, treatment

SPECIES:

Swine

ROA:

dw

DOSE:

1 jar/1,682 L (370 Imperial gal) drinking water (0.2 g/L drinking water).

1 pail/50,460 L (11,100 Imperial gal) drinking water (20 g/L drinking water).

(Provides ~297,000 I.U. penicillin/ L).

Give as sole source of drinking water for 5d.

WITHDRAWAL TIME:

Meat: 1d

USE:

Bacterial infections, systemic prevention
Bacterial infections, systemic treatment
Erysipelas, prevention/treatment
Streptococcus suis infections, treatment

SPECIES:

Turkeys

ROA:

dw

DOSE:

1 jar/1,682 L (370 Imperial gal) drinking water (0.2 g/L drinking water).

1 pail/50,460 L (11,100 Imperial gal) drinking water (20 g/L drinking water).

(Provides ~297,000 I.U. penicillin/ L).

Give as sole source of drinking water for 3-5d.

WITHDRAWAL TIME:

Meat: 1d

WT NOTES:

Do not use in birds producing eggs for food purposes.