



# NeoMed 325

## INGREDIENT(S):

neomycin sulfate

## USE:

Enteritis (bacterial), treatment

## SPECIES:

Beef Cattle, Dairy Cattle

## ROA:

dw

## DOSE:

Calves:

Individual treatment: 1 g/50 kg b.w. to daily drinking water.

Herd treatment: 100 g/5,000 kg b.w. to daily drinking water.

## WITHDRAWAL TIME:

Meat: 30d

## USE:

Bacterial infections, systemic treatment

Enteritis (bacterial), treatment

Enteritis (CRD-related), treatment

## SPECIES:

Chickens

## ROA:

dw

## DOSE:

Day-old to 2 weeks: 20 g/225 L of water OR 100 g/1,125 L of water.

2 weeks to adulthood: 40 g/225 L of water OR 200 g/1,125 L of water.

Treat for 3-5d.

## WITHDRAWAL TIME:

Meat: 7-14d

## USE:

Enteritis (bacterial), treatment

## SPECIES:

Horses

ROA:

dw

DOSE:

Individual treatment: 1 g/50 kg b.w. to daily drinking water.

Herd treatment: 100 g/5,000 kg b.w. to daily drinking water.

WT NOTES:

Not for use in horses intended for food.

USE:

Enteritis (bacterial), treatment

SPECIES:

Sheep, Swine

ROA:

dw

DOSE:

Individual treatment: 1 g/50 kg b.w. to daily drinking water.

Herd treatment: 100 g/5,000 kg b.w. to daily drinking water.

WITHDRAWAL TIME:

Meat: 14d

USE:

Enteritis (bacterial), treatment

Enteritis (CRD-related), treatment

SPECIES:

Turkeys

ROA:

dw

DOSE:

Bacterial enteritis/enteritis associated with CRD: 40 g powder/225 L water OR 200 g/1,125 L water.

WITHDRAWAL TIME:

Meat: 14d

USE:

Enteritis (Coronviral/Bluecomb), treatment

SPECIES:

Turkeys

ROA:

dw

DOSE:

Bluecomb or transmissible enteritis: 100 g/335 L water for 8d of medication as per the following schedule:

- a) 3 days, medication
- b) 2 days, no medication
- c) 3 days, medication
- d) 2 days, no medication
- e) 2 days, medication.

WITHDRAWAL TIME:

---

Meat: 14d