



Ractopamine 100

INGREDIENT(S):

ractopamine hydrochloride

USE:

Feed efficiency improvement

Weight gain increase (growth promotion)

SPECIES:

Beef Cattle

ROA:

In the feed

DOSE:

Feed continuously as sole ration to finishing cattle >400 kg b.w. for 28-42d prior to slaughter: 10-30 ppm in final feed.

Thoroughly mix 100-300 g/1,000 kg of complete beef feed (100% dry matter basis) to provide from 10-30 ppm ractopamine hydrochloride/tonne of feed.

WITHDRAWAL TIME:

Meat: 0d

WT NOTES:

Do not use in calves to be processed for veal.

Do not use in lactating dairy cattle.

USE:

Carcass leanness, increase

SPECIES:

Beef Cattle

ROA:

In the feed

DOSE:

Feed continuously as sole ration to finishing cattle >400 kg b.w. for 28-42d prior to slaughter: 20-30 ppm in final feed.

Thoroughly mix 200-300 g/1,000 kg of complete beef feed (100% dry matter basis) to provide from 20-30 ppm ractopamine hydrochloride/tonne of feed.

WITHDRAWAL TIME:

Meat: 0d

WT NOTES:

Do not use in calves to be processed for veal.

Do not use in lactating dairy cattle.