



# Tylosin 40

## INGREDIENT(S):

tylosin phosphate

## USE:

Liver abscesses, reduction of incidence

## SPECIES:

Beef Cattle

## ROA:

In the feed

## DOSE:

125 g/tonne complete diet (11 g tylosin). Feed continuously as sole ration. NOTE: Not for use in thixotropic liquid supplements.

## WITHDRAWAL TIME:

Meat: 0d

## WT NOTES:

Do not use in lactating cattle.

## USE:

Necrotic enteritis (*Clostridium perfringens*), treatment

## SPECIES:

Chickens

## ROA:

In the feed

## DOSE:

Broiler chickens: 2.27 kg (200 g tylosin)/1,000 kg complete feed. Feed as sole ration for 7d.

## WITHDRAWAL TIME:

Meat: 0d

## WT NOTES:

Do not use in laying hens.

## USE:

Porcine proliferative enteritis (ileitis), prevention/control

## SPECIES:

Swine

## ROA:

In the feed

## DOSE:

1.25 kg/tonne complete feed (110 g tylosin activity). Feed as sole ration for 21d, starting before an expected outbreak of disease.

**WITHDRAWAL TIME:**

Meat: 0d

**WT NOTES:**

No preslaughter withdrawal period is required when swine are treated with this drug at a level of 110 g tylosin per 1,000 kg of complete feed, except when used concurrently with tylosin in the drinking water.

Swine treated concurrently with this drug in complete feed and with tylosin tartrate in drinking water must not be slaughtered for use in food for at least 48 hours after the last treatment with tylosin in drinking water.

**USE:**

Swine dysentery, control/prevention

**SPECIES:**

Swine

**ROA:**

In the feed

**DOSE:**

Prevention of recurrence after treatment: 0.5 kg/tonne complete feed (44 g tylosin activity) until market weight. Feed as sole ration.

**WITHDRAWAL TIME:**

Meat: 0d

**WT NOTES:**

No preslaughter withdrawal period is required when swine are treated with this drug at a level of 110 g tylosin per 1,000 kg of complete feed, except when used concurrently with tylosin in the drinking water.

Swine treated concurrently with this drug in complete feed and with tylosin tartrate in drinking water must not be slaughtered for use in food for at least 48 hours after the last treatment with tylosin in drinking water.

**USE:**

Swine dysentery, treatment

**SPECIES:**

Swine

**ROA:**

In the feed

**DOSE:**

Use concurrently with tylosin drinking water treatment.

1.25 kg (110 g tylosin)/1,000 kg complete feed for 21d. Medicated drinking water, containing 1 g tylosin tartrate/U.S. gal (3.785 L) should be provided for first 3d of treatment, or until symptoms disappear.

**WITHDRAWAL TIME:**

Meat: 48h

**WT NOTES:**

No preslaughter withdrawal period is required when swine are treated with this drug at a level of 110 g tylosin per 1,000 kg of complete feed, except when used concurrently with tylosin in the drinking water.

Swine treated concurrently with this drug in complete feed and with tylosin tartrate in drinking water must not be slaughtered for use in food for at least 48 hours after the last treatment with tylosin in drinking water.

USE:

---

Porcine proliferative enteritis (ileitis), treatment

SPECIES:

---

Swine

ROA:

---

In the feed

DOSE:

---

Aid in treatment of PPE following treatment with tylosin drinking water treatment: Treat by way of drinking water for 7d at 1 g/12 L (83 mg/L) followed by premix in feed at a rate of 1.25 kg (110 g tylosin)/1,000 kg complete feed for 7d.

Treatment: 1.25 kg (110 g tylosin)/1,000 kg complete feed for 3 weeks.

Treatment in the presence of persistent or recurring infection: 1.25 kg (110 g tylosin)/1,000 kg complete feed for 3 weeks. Follow with 0.5 kg (44 g tylosin)/1,000 kg complete feed for 3 weeks.

WITHDRAWAL TIME:

---

Meat: 0d

WT NOTES:

---

No preslaughter withdrawal period is required when swine are treated with this drug at a level of 110 g tylosin per 1,000 kg of complete feed, except when used concurrently with tylosin in the drinking water.

Swine treated concurrently with this drug in complete feed and with tylosin tartrate in drinking water must not be slaughtered for use in food for at least 48 hours after the last treatment with tylosin in drinking water.