



Amprolium 250

INGREDIENT(S):

amprolium

USE:

Coccidiosis, control/prevention

SPECIES:

Beef Cattle, Dairy Cattle

ROA:

In the feed

DOSE:

250 mg/50 kg (5 mg/kg) b.w. daily for 21d during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard.

WITHDRAWAL TIME:

Meat: 7d

USE:

Coccidiosis, treatment

SPECIES:

Beef Cattle, Dairy Cattle

ROA:

In the feed

DOSE:

500 mg/50 kg (10 mg/kg) b.w. daily for 5d. Mix proper amount product with the amount of ration usually consumed in one day.

WITHDRAWAL TIME:

Meat: 7d

USE:

Coccidiosis, control/prevention

SPECIES:

Chickens

ROA:

In the feed

DOSE:

Broilers (death prevention due to coccidiosis): Use 0.0125% or 0.025% amprolium medicated feed as sole ration from 1d of age until birds are slaughtered.

Replacement chickens (death prevention due to coccidiosis) – floor managed layers: Heavy exposure:

0-5 weeks of age: Use 0.0125%

5-9 weeks of age: Use 0.0125%-0.008%

9-14 weeks of age: Use 0.008%

Light to moderate exposure:

0-5 weeks of age: Use 0.008%

5-9 weeks of age: Use 0.008%-0.006%

9-14 weeks of age: Use 0.006%

Replacement chickens (death prevention due to coccidiosis) – caged layers: Use 0.0125% or 0.025% amprolium medicated feed as sole ration from 1d of age until birds are placed in cages.

Mixing rates:

Amprolium level / Amount Amprolium 250 per 1,000 kg complete feed

0.006 % / 240 g

0.008% / 320 g

0.0125% / 500 g

0.025 % / 1 kg

0.05 % / 2 kg.

WT NOTES:

Do not feed to laying birds in production.

USE:

Coccidiosis, control/prevention

SPECIES:

Turkeys

ROA:

In the feed

DOSE:

Use 0.0125% amprolium (500 g/1,000 kg complete feed) as sole ration from 1d of age up to and including 16 weeks of age.

WT NOTES:

Do not feed to laying birds in production.

