



Bacitracin MD

INGREDIENT(S):

bacitracin methylene disalicylate

USE:

Necrotic enteritis (*Clostridium perfringens*), control/prevention

SPECIES:

Chickens

ROA:

In the feed

DOSE:

Broilers: 0.5 kg/999.5 kg complete feed (55 mg/kg). Feed continuously to market weight.

WITHDRAWAL TIME:

Meat: 0d

WT NOTES:

This product has a zero day withdrawal period for eggs.

USE:

Early mortality, reduction

SPECIES:

Chickens

ROA:

In the feed

DOSE:

Broilers (due to diminished feed consumption and chilling): 1 kg/999 kg complete feed (110 mg/kg). Feed to 1 week of age.

WITHDRAWAL TIME:

Meat: 0d

WT NOTES:

This product has a zero day withdrawal period for eggs.

USE:

Necrotic enteritis (*Clostridium perfringens*), treatment

SPECIES:

Chickens

ROA:

In the feed

DOSE:

Laying hens and Broilers: Start medication at first clinical signs of disease. Mix 1 kg/999 kg complete feed (providing 110 mg/kg). Feed continuously for 7d.

WITHDRAWAL TIME:

Meat: 0d

WT NOTES:

This product has a zero day withdrawal period for eggs.

USE:

Enteritis (*Clostridium perfringens*), prevention

SPECIES:

Swine

ROA:

In the feed

DOSE:

Pregnant and lactating sows or gilts: Mix 2.5 kg/997.5 kg complete feed (275 mg/kg). Feed continuously for 2 1/2 weeks before farrowing until 3 weeks after farrowing.

WITHDRAWAL TIME:

Meat: 0d