



# Bacitracin MD

## INGREDIENT(S):

bacitracin methylene disalicylate

## USE:

Necrotic enteritis (*Clostridium perfringens*), control/prevention

## SPECIES:

Chickens

## ROA:

In the feed

## DOSE:

Broilers: 0.5 kg/999.5 kg complete feed (55 mg/kg). Feed continuously to market weight.

## WITHDRAWAL TIME:

Meat: 0d

## WT NOTES:

This product has a zero day withdrawal period for eggs.

## USE:

Early mortality, reduction

## SPECIES:

Chickens

## ROA:

In the feed

## DOSE:

Broilers (due to diminished feed consumption and chilling): 1 kg/999 kg complete feed (110 mg/kg). Feed to 1 week of age.

## WITHDRAWAL TIME:

Meat: 0d

## WT NOTES:

This product has a zero day withdrawal period for eggs.

## USE:

Necrotic enteritis (*Clostridium perfringens*), treatment

## SPECIES:

Chickens

## ROA:

In the feed

## DOSE:

Laying hens and Broilers: Start medication at first clinical signs of disease. Mix 1 kg/999 kg complete feed (providing 110 mg/kg). Feed continuously for 7d.

**WITHDRAWAL TIME:**

---

Meat: 0d

**WT NOTES:**

---

This product has a zero day withdrawal period for eggs.

**USE:**

---

Enteritis (*Clostridium perfringens*), prevention

**SPECIES:**

---

Swine

**ROA:**

---

In the feed

**DOSE:**

---

Pregnant and lactating sows or gilts: Mix 2.5 kg/997.5 kg complete feed (275 mg/kg). Feed continuously for 2 1/2 weeks before farrowing until 3 weeks after farrowing.

**WITHDRAWAL TIME:**

---

Meat: 0d