



# Chlor 50 Granular Premix

**INGREDIENT(S):**

chlortetracycline

**USE:**

Enteritis (bacterial), prevention

**SPECIES:**

Beef Cattle, Dairy Cattle

**ROA:**

In the feed

**DOSE:**

Calves up to 136 kg: Feed continuously during periods of early growth.

In the feed: Add 0.5 kg/tonne [0.0055% (55 mg/kg)].

Milk replacer: Add 0.5 kg/tonne [0.0055% (55 mg/kg)].

**WITHDRAWAL TIME:**

Meat: 5d

**WT NOTES:**

Do not administer to lactating dairy cattle.

**USE:**

Footrot, prevention

**SPECIES:**

Beef Cattle, Dairy Cattle

**ROA:**

In the feed

**DOSE:**

Add 200 g/tonne (0.22 mg/kg/day b.w.) or add 640 g/tonne supplement and feed 1 kg/head/day (70 mg/head/day).

Feed continuously during exposure period.

**WITHDRAWAL TIME:**

Meat: 5d

**WT NOTES:**

Do not administer to lactating dairy cattle.

**USE:**

Enterotoxemia (*Clostridium perfringens*), reduction of losses

**SPECIES:**

Sheep

ROA:

In the feed

DOSE:

Feedlot lambs: 0.0022% (22 mg/kg) complete feed. Feed continuously as sole ration while lambs are in feedlot.

WITHDRAWAL TIME:

Meat: 4d

USE:

Enteritis (bacterial), treatment

SPECIES:

Swine

ROA:

In the feed

DOSE:

Add 1 kg/tonne [0.011% (110 mg/kg)]. Feed continuously as sole ration following the appearance of symptoms and until 3d after symptoms disappear.

Prestarter feeds and feeds to be fed for up to and including 6 weeks of age are required to contain 0.022%.

WITHDRAWAL TIME:

Meat: 7d

USE:

Enteritis (bacterial), prevention

SPECIES:

Swine

ROA:

In the feed

DOSE:

0.5 kg/tonne [0.0055% (55 mg/kg)]. Feed continuously as sole ration during early growth period up to 32 kg (70 lb) b.w.

Prestarter feeds and feeds to be fed for up to and including 6 weeks of age are required to contain 0.022%.

WITHDRAWAL TIME:

Meat: 7d

USE:

Hexamitiasis, control/prevention

SPECIES:

Turkeys

ROA:

In the feed

**DOSE:**

0.0055% (55 mg/kg) complete feed: Feed as sole ration for as long as birds are expected to be exposed to infection.

**WITHDRAWAL TIME:**

Meat: 7

**USE:**

Hexamitiasis, treatment

**SPECIES:**

Turkeys

**ROA:**

In the feed

**DOSE:**

0.011% (110 mg/kg) complete feed: Feed as sole ration when symptoms appear and continue for a few days after symptoms disappear.

**WITHDRAWAL TIME:**

Meat: 7d

**USE:**

Synovitis (infectious – *Mycoplasma synoviae*), treatment

**SPECIES:**

Turkeys

**ROA:**

In the feed

**DOSE:**

0.022% (220 mg/kg) complete feed: Feed as sole ration when symptoms appear and continue for at least 2 weeks after symptoms disappear.

**WITHDRAWAL TIME:**

Meat: 7d