



Lincomycin 110 G

INGREDIENT(S):

lincomycin hydrochloride

USE:

Swine dysentery (*Brachyspira* spp), control/prevention

SPECIES:

Swine

ROA:

In the feed

DOSE:

Growing swine: 0.4 kg/tonne of complete feed (44 g of lincomycin base/tonne). Feed as sole ration for 4 weeks.
Repeat prn.

WITHDRAWAL TIME:

Meat: 0d

WT NOTES:

No drug withdrawal period is required before slaughter for swine medicated with lincomycin at the recommended level of 44 or 110 grams of lincomycin base per tonne.
Swine treated at the recommended level of 220 grams of lincomycin per tonne must not be slaughtered for use in food for at least 24 hours after the latest treatment with this drug.

USE:

Swine dysentery (*Brachyspira* spp), treatment

SPECIES:

Swine

ROA:

In the feed

DOSE:

Growing swine: 1 kg/tonne of complete feed (110 g of lincomycin base/tonne). Feed as sole ration for 21d, or until signs of disease start to disappear, then feed at preventative level.

WITHDRAWAL TIME:

Meat: 24h

WT NOTES:

No drug withdrawal period is required before slaughter for swine medicated with lincomycin at the recommended level of 44 or 110 grams of lincomycin base per tonne.
Swine treated at the recommended level of 220 grams of lincomycin per tonne must not be slaughtered for use in food for at least 24 hours after the latest treatment with this drug.

USE:

Mycoplasmal pneumonia, reduction of severity

SPECIES:

Swine

ROA:

In the feed

DOSE:

Growing swine: 2 kg/tonne of complete feed (220 g lincomycin base/tonne). Feed as sole ration for 21d.

WITHDRAWAL TIME:

Meat: 24h

WT NOTES:

No drug withdrawal period is required before slaughter for swine medicated with lincomycin at the recommended level of 44 or 110 grams of lincomycin base per tonne.

Swine treated at the recommended level of 220 grams of lincomycin per tonne must not be slaughtered for use in food for at least 24 hours after the latest treatment with this drug.

USE:

Porcine proliferative enteritis (ileitis), reduction of severity

SPECIES:

Swine

ROA:

In the feed

DOSE:

1 kg/tonne complete feed (110 g of lincomycin base/tonne). Feed as sole ration for 21d.

WITHDRAWAL TIME:

Meat: 0d

WT NOTES:

No drug withdrawal period is required before slaughter for swine medicated with lincomycin at the recommended level of 44 or 110 grams of lincomycin base per tonne.

Swine treated at the recommended level of 220 grams of lincomycin per tonne must not be slaughtered for use in food for at least 24 hours after the latest treatment with this drug.