



# Lincomycin 110 G

## INGREDIENT(S):

lincomycin hydrochloride

## USE:

Swine dysentery (*Brachyspira* spp), control/prevention

## SPECIES:

Swine

## ROA:

In the feed

## DOSE:

Growing swine: 0.4 kg/tonne of complete feed (44 g of lincomycin base/tonne). Feed as sole ration for 4 weeks. Repeat prn.

## WITHDRAWAL TIME:

Meat: 0d

## WT NOTES:

No drug withdrawal period is required before slaughter for swine medicated with lincomycin at the recommended level of 44 or 110 grams of lincomycin base per tonne.

Swine treated at the recommended level of 220 grams of lincomycin per tonne must not be slaughtered for use in food for at least 24 hours after the latest treatment with this drug.

## USE:

Swine dysentery (*Brachyspira* spp), treatment

## SPECIES:

Swine

## ROA:

In the feed

## DOSE:

Growing swine: 1 kg/tonne of complete feed (110 g of lincomycin base/tonne). Feed as sole ration for 21d, or until signs of disease start to disappear, then feed at preventative level.

## WITHDRAWAL TIME:

Meat: 24h

## WT NOTES:

No drug withdrawal period is required before slaughter for swine medicated with lincomycin at the recommended level of 44 or 110 grams of lincomycin base per tonne.

Swine treated at the recommended level of 220 grams of lincomycin per tonne must not be slaughtered for use in food for at least 24 hours after the latest treatment with this drug.

## USE:

Mycoplasmal pneumonia, reduction of severity

**SPECIES:**

Swine

**ROA:**

In the feed

**DOSE:**

Growing swine: 2 kg/tonne of complete feed (220 g lincomycin base/tonne). Feed as sole ration for 21d.

**WITHDRAWAL TIME:**

Meat: 24h

**WT NOTES:**

No drug withdrawal period is required before slaughter for swine medicated with lincomycin at the recommended level of 44 or 110 grams of lincomycin base per tonne.

Swine treated at the recommended level of 220 grams of lincomycin per tonne must not be slaughtered for use in food for at least 24 hours after the latest treatment with this drug.

**USE:**

Porcine proliferative enteritis (ileitis), reduction of severity

**SPECIES:**

Swine

**ROA:**

In the feed

**DOSE:**

1 kg/tonne complete feed (110 g of lincomycin base/tonne). Feed as sole ration for 21d.

**WITHDRAWAL TIME:**

Meat: 0d

**WT NOTES:**

No drug withdrawal period is required before slaughter for swine medicated with lincomycin at the recommended level of 44 or 110 grams of lincomycin base per tonne.

Swine treated at the recommended level of 220 grams of lincomycin per tonne must not be slaughtered for use in food for at least 24 hours after the latest treatment with this drug.