

Lincomycin Spectinomycin 100 Soluble Powder

INGREDIENT(S):
lincomycin hydrochloride, spectinomycin
USE:
Chronic respiratory disease/infectious sinusitis, control/prevention
SPECIES:
Chickens
ROA:
dw
DOSE:
1 jar/120 L drinking water (833 mg antibiotic activity/L).
Automatic medicators: Set to deliver 7.8 mL stock solution/L drinking water. Stock solution: 1 jar/0.9 L.
To prevent losses due to CRD, administer for first 3-5d of life and again for 1-2d at ~4 weeks of age.
WITHDRAWAL TIME: Meat: 3d
Wiedt. 3d
WT NOTES:
This product must not be used in laying birds.
USE:
Chronic respiratory disease/infectious sinusitis, treatment
SPECIES:
Chickens
ROA:
dw
DOSE.
DOSE: 1 jar/120 L drinking water (833 mg antibiotic activity/L).
Thus 120 2 drinking water (055 mg andorotic activity/2).
Automatic medicators: Set to deliver 7.8 mL stock solution/L drinking water. Stock solution: 1 jar/0.9 L.
Use as sole source of drinking water for 5-7d.
WITHDRAWAL TIME:
Meat: 3d
WT NOTES:
This product must not be used in laying birds.
USE:

Swine dysentery, treatment
SPECIES:
Swine
ROA:
dw
DOSE:
1 jar/1,500 L drinking water (67 mg antibiotic activity/L).
Automatic medicators: Set to deliver 7.8 mL stock solution/L drinking water. Stock solution: 1 jar/11.7 L.
Use as sole source of drinking water for 4-7d.
WITHOR AWAI TIME

Meat: 5d