



# Neo-Tetramed

**INGREDIENT(S):**

neomycin sulfate, tetracycline

**USE:**

Enteritis (bacterial), treatment

**SPECIES:**

Beef Cattle, Dairy Cattle

**ROA:**

dw, mr

**DOSE:**

Individual treatment of calves: 2 g /60 mL of water or milk replacer/45 kg b.w. as a drench.

Herd treatment of calves: 2 g/4.5 L of water OR 100 g (1 pouch)/225 L water for 4-5d.

**WITHDRAWAL TIME:**

Meat: 30d

**USE:**

Respiratory disease (bacterial), control/treatment

**SPECIES:**

Beef Cattle, Dairy Cattle

**ROA:**

dw, mr

**DOSE:**

Individual treatment of calves: 4 g /60 mL of water or milk substitute per 45 kg b.w. as a drench.

Herd treatment of calves: 4 g/4.5 L of water OR 200 g (2 pouches)/225 L drinking water for 4-5d.

**WITHDRAWAL TIME:**

Meat: 30d

**USE:**

Chronic respiratory disease/infectious sinusitis, treatment

Enteritis (bacterial), treatment

Synovitis (infectious – Mycoplasma synoviae), treatment

**SPECIES:**

Chickens

**ROA:**

dw

**DOSE:**

400-800 g/900 L drinking water for 3-5d.

**WITHDRAWAL TIME:**

Meat: 7d

**WT NOTES:**

This product must not be used in laying birds.

**USE:**

Enteritis (bacterial), treatment

**SPECIES:**

Swine

**ROA:**

dw

**DOSE:**

Swine: 2 g/170 mL of water per 45 kg b.w. as a drench OR 100 g/225 L drinking water for 4-5 d.

Newborn piglets: 4 g/60 mL water and give 5 mL/piglet q12h for 4-5d.

**WITHDRAWAL TIME:**

Meat: 14d

**USE:**

Respiratory disease (bacterial), control/treatment

**SPECIES:**

Swine

**ROA:**

dw

**DOSE:**

4 g/170 mL of water per 45 kg b.w. OR 200 g/225 L of drinking water and give for 4-5d.

**WITHDRAWAL TIME:**

Meat: 14d

**USE:**

Chronic respiratory disease/infectious sinusitis, treatment

Enteritis (bacterial), treatment

Enteritis (Coronviral/Bluecomb), treatment

Synovitis (infectious – Mycoplasma synoviae), treatment

**SPECIES:**

Turkeys

**ROA:**

dw

**DOSE:**

400-800 g/900 L drinking water for 3-5d.

**WITHDRAWAL TIME:**

---

Meat: 14d

**WT NOTES:**

---

This product must not be used in laying birds.