



Neo-Tetramed

INGREDIENT(S):

neomycin sulfate, tetracycline

USE:

Enteritis (bacterial), treatment

SPECIES:

Beef Cattle, Dairy Cattle

ROA:

dw, mr

DOSE:

Individual treatment of calves: 2 g /60 mL of water or milk replacer/45 kg b.w. as a drench.

Herd treatment of calves: 2 g/4.5 L of water OR 100 g (1 pouch)/225 L water for 4-5d.

WITHDRAWAL TIME:

Meat: 30d

USE:

Respiratory disease (bacterial), control/treatment

SPECIES:

Beef Cattle, Dairy Cattle

ROA:

dw, mr

DOSE:

Individual treatment of calves: 4 g /60 mL of water or milk substitute per 45 kg b.w. as a drench.

Herd treatment of calves: 4 g/4.5 L of water OR 200 g (2 pouches)/225 L drinking water for 4-5d.

WITHDRAWAL TIME:

Meat: 30d

USE:

Chronic respiratory disease/infectious sinusitis, treatment

Enteritis (bacterial), treatment

Synovitis (infectious – Mycoplasma synoviae), treatment

SPECIES:

Chickens

ROA:

dw

DOSE:

400-800 g/900 L drinking water for 3-5d.

WITHDRAWAL TIME:

Meat: 7d

WT NOTES:

This product must not be used in laying birds.

USE:

Enteritis (bacterial), treatment

SPECIES:

Swine

ROA:

dw

DOSE:

Swine: 2 g/170 mL of water per 45 kg b.w. as a drench OR 100 g/225 L drinking water for 4-5 d.

Newborn piglets: 4 g/60 mL water and give 5 mL/piglet q12h for 4-5d.

WITHDRAWAL TIME:

Meat: 14d

USE:

Respiratory disease (bacterial), control/treatment

SPECIES:

Swine

ROA:

dw

DOSE:

4 g/170 mL of water per 45 kg b.w. OR 200 g/225 L of drinking water and give for 4-5d.

WITHDRAWAL TIME:

Meat: 14d

USE:

Chronic respiratory disease/infectious sinusitis, treatment

Enteritis (bacterial), treatment

Enteritis (Coronviral/Bluecomb), treatment

Synovitis (infectious – Mycoplasma synoviae), treatment

SPECIES:

Turkeys

ROA:

dw

DOSE:

400-800 g/900 L drinking water for 3-5d.

WITHDRAWAL TIME:

Meat: 14d

WT NOTES:

This product must not be used in laying birds.