



Oxytetracycline 200 Granular Premix

INGREDIENT(S):

oxytetracycline

USE:

Enteritis (bacterial), prevention

SPECIES:

Beef Cattle

ROA:

In the feed

DOSE:

Calves: 0.125 kg/tonne. Feed continuously during period of early growth (up to 136 kg b.w.).

WITHDRAWAL TIME:

Meat: 5d

USE:

Bloat, control/prevention

SPECIES:

Beef Cattle

ROA:

In the feed

DOSE:

0.125 kg/tonne complete feed, fed at rate of 1.36 kg/head/day.

Feed continuously so each animal receives 75 mg/day.

WITHDRAWAL TIME:

Meat: 5d

USE:

Enteritis (bacterial), prevention

SPECIES:

Dairy Cattle

ROA:

In the feed

DOSE:

Calves: 0.125 kg/tonne. Feed continuously during period of early growth (up to 136 kg b.w.).

WITHDRAWAL TIME:

Meat: 5d

WT NOTES:

Do not administer to lactating dairy cattle.

USE:

Enteritis (bacterial), prevention

SPECIES:

Sheep

ROA:

In the feed

DOSE:

0.25 kg/tonne complete feed. Feed as sole creep feed during suckling period.

WITHDRAWAL TIME:

Meat: 4d

USE:

Enterotoxemia (*Clostridium perfringens*), reduction of losses

SPECIES:

Sheep

ROA:

In the feed

DOSE:

0.05 kg/tonne complete feed. Feed as sole ration while lambs are in feed lot.

WITHDRAWAL TIME:

Meat: 4d

USE:

Enteritis (bacterial), prevention

SPECIES:

Swine

ROA:

In the feed

DOSE:

Piglets <6 weeks of age: 0.5 kg/tonne complete feed. Feed as sole ration.

Piglets ≥6 weeks of age up to 27-32 kg b.w.: 0.125 kg/tonne complete feed. Feed as sole ration.

WITHDRAWAL TIME:

Meat: 7d

USE:

Enteritis (bacterial), treatment

SPECIES:

Swine

ROA:

In the feed

DOSE:

Piglets <=6 weeks of age: 0.5 kg/tonne complete feed. Feed as sole ration when symptoms appear and until 3d asymptomatic.

Piglets >6 weeks of age: 0.25 kg/tonne complete feed. Feed as above.

WITHDRAWAL TIME:

Meat: 7d

USE:

Abortion (Leptospirosis), control/treatment

SPECIES:

Swine

ROA:

In the feed

DOSE:

1.25 kg/tonne complete feed.

Feed once diagnosis has been made as sole ration to pregnant and affected sows for 2 weeks.

WITHDRAWAL TIME:

Meat: 7d

USE:

Chronic respiratory disease/infectious sinusitis, control/prevention
Synovitis (infectious – Mycoplasma synoviae), control/prevention

SPECIES:

Turkeys

ROA:

In the feed

DOSE:

0.25 kg/tonne. Feed as sole ration for duration of exposure.

WITHDRAWAL TIME:

Meat: 7d

USE:

Synovitis (infectious – Mycoplasma synoviae), treatment

SPECIES:

Turkeys

ROA:

In the feed

DOSE:

0.5 kg/tonne.

Feed as sole ration when symptoms appear and continue until 2 weeks after they disappear.

WITHDRAWAL TIME:

Meat: 7d
