



# Pot-Pen® 15 000 000 000 I.U.

**INGREDIENT(S):**

penicillin G potassium

**USE:**

Necrotic enteritis (*Clostridium perfringens*), control/prevention

**SPECIES:**

Chickens

**ROA:**

dw

**DOSE:**

1 jar/1,682 L (370 Imperial gal) drinking water (0.2 g/L drinking water).

1 pail/50,460 L (11,100 Imperial gal) drinking water (20 g/L drinking water).

(Provides ~297,000 I.U. penicillin/ L).

Give as sole source of drinking water for 5d.

**WITHDRAWAL TIME:**

Meat: 1d

**USE:**

*Streptococcus suis* infections, treatment

**SPECIES:**

Swine

**ROA:**

dw

**DOSE:**

1 jar/1,682 L (370 Imperial gal) drinking water (0.2 g/L drinking water).

1 pail/50,460 L (11,100 Imperial gal) drinking water (20 g/L drinking water).

(Provides ~297,000 I.U. penicillin/ L).

Give as sole source of drinking water for 5d.

**WITHDRAWAL TIME:**

Meat: 1d

**USE:**

Bacterial infections, systemic prevention  
Bacterial infections, systemic treatment  
Erysipelas, prevention/treatment  
Streptococcus suis infections, treatment

SPECIES:

---

Turkeys

ROA:

---

dw

DOSE:

---

1 jar/1,682 L (370 Imperial gal) drinking water (0.2 g/L drinking water).

1 pail/50,460 L (11,100 Imperial gal) drinking water (20 g/L drinking water).

(Provides ~297,000 I.U. penicillin/ L).

Give as sole source of drinking water for 3-5d.

WITHDRAWAL TIME:

---

Meat: 1d

WT NOTES:

---

Do not use in birds producing eggs for food purposes.