



Ractopamine 4



Ractopamine 4

INGREDIENT(S):

ractopamine hydrochloride

USE:

Feed efficiency improvement
Weight gain increase (growth promotion)

SPECIES:

Beef Cattle

ROA:

In the feed

DOSE:

10 ppm: 2.5 kg (1/2 bag)/1,000 kg feed.

20 ppm: 5 kg (1 bag)/1,000 kg feed.

30 ppm: 7.5 kg (1 1/2 bags)/1,000 kg feed.

Finishing cattle >400 kg b.w.: Feed continuously as sole ration for 28-42d prior to slaughter.

WITHDRAWAL TIME:

Meat: 0d

WT NOTES:

Do not use in calves to be processed for veal.
Do not use in lactating dairy cattle.

USE:

Carcass leanness, increase

SPECIES:

Beef Cattle

ROA:

In the feed

DOSE:

20 ppm: 5 kg (1 bag)/1,000 kg feed.

30 ppm: 7.5 kg (1 1/2 bags)/1,000 kg feed.

Finishing cattle >400 kg b.w.: Feed continuously as sole ration for 28-42d prior to slaughter.

WITHDRAWAL TIME:

Meat: 0d

WT NOTES:

Do not use in calves to be processed for veal.

Do not use in lactating dairy cattle.