



Terramycin®-100 Premix

INGREDIENT(S):

oxytetracycline

USE:

Bloat, control/prevention

SPECIES:

Beef Cattle

ROA:

In the feed

DOSE:

75 mg oxytetracycline/day.

Mix 55 g/tonne (0.25 kg product) of complete feed. Feed 1.36 kg/head/day (75 mg conc.).

WITHDRAWAL TIME:

Meat: 5d

USE:

Enteritis (bacterial), prevention

SPECIES:

Beef Cattle, Dairy Cattle

ROA:

In the feed

DOSE:

Calves: Feed continuously during period of early growth (up to 136 kg b.w.).

In the feed: 0.25 kg/tonne.

WITHDRAWAL TIME:

Meat: 5d

WT NOTES:

Do not administer to lactating dairy cattle.

USE:

Enteritis (bacterial), prevention

SPECIES:

Sheep

ROA:

In the feed

DOSE:

0.5 kg/tonne of complete feed. Feed as sole ration during the suckling period.

WITHDRAWAL TIME:

Meat: 4d

USE:

Enterotoxemia (Clostridium perfringens), reduction of losses

SPECIES:

Sheep

ROA:

In the feed

DOSE:

0.1 kg/tonne of complete feed. Feed as sole ration while lambs are in feedlot.

WITHDRAWAL TIME:

Meat: 4d

USE:

Enteritis (bacterial), prevention

SPECIES:

Swine

ROA:

In the feed

DOSE:

Piglets <6 weeks of age: 1 kg/tonne complete feed. Feed as sole ration.

Piglets 6 weeks to 27-31 kg b.w.: 0.25 kg/tonne complete feed. Feed as sole ration.

WITHDRAWAL TIME:

Meat: 7d

USE:

Enteritis (bacterial), treatment

SPECIES:

Swine

ROA:

In the feed

DOSE:

Piglets <6 weeks of age: 1 kg/tonne complete feed. Feed as sole ration following appearance of symptoms and for 3d after they disappear.

Piglets 6 weeks to 27-31 kg b.w.: 0.5 kg/tonne complete feed. Feed as above.

WITHDRAWAL TIME:

Meat: 7d

USE:

Abortion (Leptospirosis), control/treatment

SPECIES:

Swine

ROA:

In the feed

DOSE:

2.5 kg/tonne complete feed. Feed as sole ration for 2 weeks to pregnant and affected sows.

WITHDRAWAL TIME:

Meat: 7d

USE:

Enteritis (non-specific), periods of

SPECIES:

Turkeys

ROA:

In the feed

DOSE:

As an aid in stimulating appetite and maintaining weight gains in the presence of non-specific enteritis: 0.50 kg/tonne complete feed. Feed as sole ration from onset of disease for at least 1 month or until a few days after symptoms disappear.

WITHDRAWAL TIME:

Meat: 7d

USE:

Chronic respiratory disease/infectious sinusitis, control/prevention

SPECIES:

Turkeys

ROA:

In the feed

DOSE:

As an aid in stimulating appetite and maintaining weight gains when chickens are expected to be exposed to mild attacks of CRD: 0.25 kg/tonne. Feed as sole ration for as long as birds are expected to be exposed to infection.

WITHDRAWAL TIME:

Meat: 7d

USE:

Enteritis (bacterial), prevention

SPECIES:

Turkeys

ROA:

In the feed

DOSE:

0.25 kg/tonne. Feed as sole ration for duration of exposure.

WITHDRAWAL TIME:

Meat: 7d

USE:

Synovitis (infectious – *Mycoplasma synoviae*), control/prevention

SPECIES:

Turkeys

ROA:

In the feed

DOSE:

0.5 kg/tonne. Feed as sole ration for duration of exposure.

WITHDRAWAL TIME:

Meat: 7d

USE:

Synovitis (infectious – *Mycoplasma synoviae*), treatment

SPECIES:

Turkeys

ROA:

In the feed

DOSE:

1 kg/tonne. Feed as sole ration when symptoms appear and continue until 2 weeks after symptoms disappear.

WITHDRAWAL TIME:

Meat: 7d