



Tetramed 1000

INGREDIENT(S):

tetracycline

USE:

Enteritis (bacterial), treatment

SPECIES:

Beef Cattle, Dairy Cattle

ROA:

dw, Oral

DOSE:

0.25 g/50 kg b.w. q12h, as a drench in water OR 125 g/1,125 L of water for 3-5d.

WITHDRAWAL TIME:

Meat: 5d

WT NOTES:

Not for use in lactating dairy animals.

USE:

Respiratory disease (bacterial), control/treatment

SPECIES:

Beef Cattle, Dairy Cattle

ROA:

dw, Oral

DOSE:

0.5 g/50 kg b.w. q12h, as a drench in water OR 250 g/1,125 L of water for 3-5d.

WITHDRAWAL TIME:

Meat: 5d

WT NOTES:

Not for use in lactating dairy animals.

USE:

Chronic respiratory disease/infectious sinusitis, treatment

Enteritis (bacterial), treatment

Fowl cholera, treatment

Hexamitiasis, treatment

SPECIES:

Chickens

ROA:

dw

DOSE:

125 g/2,500 L water for 3-5d.

Automatic proportioner: Set apparatus to distribute 15 mL/4 L drinking water (1/2 oz/US gal).

Prepare stock solution by dissolving 125 g/9 L water. This will medicate 2,460 L drinking water.

WITHDRAWAL TIME:

Meat: 5d

WT NOTES:

Do not feed to laying birds.

USE:

Enteritis (bacterial), treatment

Respiratory disease (bacterial), control/treatment

SPECIES:

Sheep

ROA:

Oral

DOSE:

0.25 g/25 kg b.w. q12h as a drench in water for 4-5d.

WITHDRAWAL TIME:

Meat: 5d

USE:

Enteritis (bacterial), treatment

Respiratory disease (bacterial), control/treatment

SPECIES:

Swine

ROA:

dw, Oral

DOSE:

0.25 g/25 kg b.w. q12h as a drench using a dose syringe or 0.5 g/4 L of drinking water or 125 g/1,000 L of drinking water for 4 or 5d.

Automatic proportioner: Set the apparatus to distribute 30 mL/4 L drinking water (1 oz/US gal). Prepare stock solution by dissolving 125 g/7.5 L of water. This will medicate 1,000 L of drinking water.

WITHDRAWAL TIME:

Meat: 5d

USE:

Chronic respiratory disease/infectious sinusitis, treatment

Enteritis (bacterial), treatment

Enteritis (Coronaviral/Bluecomb), treatment

Fowl cholera, treatment

Hexamitiasis, treatment

SPECIES:

Turkeys

ROA:

dw

DOSE:

125 g/2,500 L water for 3-5d.

Automatic proportioner: Set apparatus to distribute 15 mL/4 L drinking water (1/2 oz/US gal).

Prepare stock solution by dissolving 125 g/9 L water. This will medicate 2,460 L drinking water.

WITHDRAWAL TIME:

Meat: 5d

WT NOTES:

Do not feed to laying birds.