



# Tetramed 250

**INGREDIENT(S):**

tetracycline

**USE:**

Enteritis (bacterial), treatment

**SPECIES:**

Beef Cattle, Dairy Cattle

**ROA:**

dw, Oral

**DOSE:**

1 g/50 kg b.w. q12h as a drench in water OR 100 g (1 pouch)/225 L of water for 3-5d.

**WITHDRAWAL TIME:**

Meat: 5d

**WT NOTES:**

Do not use in lactating dairy cattle.

**USE:**

Respiratory disease (bacterial), control/treatment

**SPECIES:**

Beef Cattle, Dairy Cattle

**ROA:**

dw, Oral

**DOSE:**

2 g/50 kg b.w. q12h as a drench in water OR 200 g (2 pouches)/225 L of water for 3-5d.

**WITHDRAWAL TIME:**

Meat: 5d

**WT NOTES:**

Do not use in lactating dairy cattle.

**USE:**

Chronic respiratory disease/infectious sinusitis, treatment

Enteritis (bacterial), treatment

Fowl cholera, treatment

**SPECIES:**

Chickens

**ROA:**

dw

**DOSE:**

1 g/5 L of drinking water OR 100 g (1 pouch)/500 L of water for 3-5d.

Automatic proportioner: Set to distribute 30 mL/4 L drinking water (1 oz/gal US). Prepare stock solution by dissolving 100 g/3.8 L water to treat 500 L drinking water OR set to distribute 15 mL/4 L water (1/2 oz/gal). Prepare stock solution by dissolving 400 g/7.5 L water to medicate 2,000 L drinking water.

**WITHDRAWAL TIME:**

Meat: 5d

**WT NOTES:**

Do not feed to laying birds.

**USE:**

Enteritis (bacterial), treatment  
Respiratory disease (bacterial), control/treatment

**SPECIES:**

Sheep

**ROA:**

Oral

**DOSE:**

1 g/25 kg b.w. q12h as a drench in water for 4-5d.

**WITHDRAWAL TIME:**

Meat: 5d

**USE:**

Enteritis (bacterial), treatment  
Respiratory disease (bacterial), control/treatment

**SPECIES:**

Swine

**ROA:**

dw, Oral

**DOSE:**

1 g/25 kg b.w. q12h as a drench OR 2 g/4 L drinking water, OR 100 g (1 pouch)/200 L drinking water OR 400 g/800 L water for 4-5d.

Automatic proportioner: Set to distribute 30 mL/4 L drinking water (1 oz/gal US). Prepare stock solution by dissolving 200 g in 3 L water OR 400 g in 6 L water to treat 400 L and 800 L drinking water.

**WITHDRAWAL TIME:**

Meat: 5d

**USE:**

Chronic respiratory disease/infectious sinusitis, treatment  
Enteritis (bacterial), treatment  
Enteritis (Coronaviral/Bluecomb), treatment  
Fowl cholera, treatment

Hexamitiasis, treatment

SPECIES:

---

Turkeys

ROA:

---

dw

DOSE:

---

1 g/5 L of drinking water OR 100 g (1 pouch)/500 L of water for 3-5d.

Automatic proportioner: Set to distribute 30 mL/4 L drinking water (1 oz/gal US). Prepare stock solution by dissolving 100 g/3.8 L water to treat 500 L drinking water OR set to distribute 15 mL/4 L water (1/2 oz/gal). Prepare stock solution by dissolving 400 g/7.5 L water to medicate 2,000 L drinking water.

WITHDRAWAL TIME:

---

Meat: 5d

WT NOTES:

---

Do not feed to laying birds.