



## Tiamulin 1.78% Premix

### INGREDIENT(S):

tiamulin

### USE:

Swine dysentery (*Brachyspira* spp), control/prevention

### SPECIES:

Swine

### ROA:

In the feed

### DOSE:

1.75 kg/tonne of complete feed (31.2 g/tonne). Feed continuously as sole ration following treatment, on premises with a history of swine dysentery but where signs of disease have not yet occurred.

### WITHDRAWAL TIME:

Meat: 2d

### WT NOTES:

Do not slaughter swine for use in food for at least 2 days after the latest treatment with this drug at 31.2 g/tonne.  
Do not slaughter swine for use in food for at least 7 days after the latest treatment with this drug at 178.1 g/tonne.

### USE:

Swine dysentery (*Brachyspira* spp), treatment

### SPECIES:

Swine

### ROA:

In the feed

### DOSE:

10 kg/tonne of complete feed (178.1 g/tonne). Feed as sole ration 14d.

### WITHDRAWAL TIME:

Meat: 7d

### WT NOTES:

Do not slaughter swine for use in food for at least 2 days after the latest treatment with this drug at 31.2 g/tonne.  
Do not slaughter swine for use in food for at least 7 days after the latest treatment with this drug at 178.1 g/tonne.

