



# Tiamulin 1.78% Premix

---

**INGREDIENT(S):**

tiamulin

**USE:**

Swine dysentery (Brachyspira spp), control/prevention

**SPECIES:**

Swine

**ROA:**

In the feed

**DOSE:**

1.75 kg/tonne of complete feed (31.2 g/tonne). Feed continuously as sole ration following treatment, on premises with a history of swine dysentery but where signs of disease have not yet occurred.

**WITHDRAWAL TIME:**

Meat: 2d

**WT NOTES:**

Do not slaughter swine for use in food for at least 2 days after the latest treatment with this drug at 31.2 g/tonne.

Do not slaughter swine for use in food for at least 7 days after the latest treatment with this drug at 178.1 g/tonne.

**USE:**

Swine dysentery (Brachyspira spp), treatment

**SPECIES:**

Swine

**ROA:**

In the feed

**DOSE:**

10 kg/tonne of complete feed (178.1 g/tonne). Feed as sole ration 14d.

**WITHDRAWAL TIME:**

Meat: 7d

**WT NOTES:**

Do not slaughter swine for use in food for at least 2 days after the latest treatment with this drug at 31.2 g/tonne.

Do not slaughter swine for use in food for at least 7 days after the latest treatment with this drug at 178.1 g/tonne.

