



# Tiamulin HF 10%

## INGREDIENT(S):

tiamulin

## USE:

Porcine colonic spirochaetosis (colitis) (*Brachyspira pilosicoli*), treatment  
Swine dysentery (*Brachyspira* spp), treatment

## SPECIES:

Swine

## ROA:

In the feed

## DOSE:

2.2 kg/tonne feed. Feed as sole ration for 14d.

## WITHDRAWAL TIME:

Meat: 7d

## USE:

Porcine proliferative enteritis (PPE, ileitis – *Lawsonia intracellularis*), treatment

## SPECIES:

Swine

## ROA:

In the feed

## DOSE:

1.5 kg/tonne feed. Feed as sole ration for 14d.

## WITHDRAWAL TIME:

Meat: 7d

## USE:

Pneumonia (enzootic – *Mycoplasma hyopneumoniae*), treatment

## SPECIES:

Swine

## ROA:

In the feed

## DOSE:

1.65 kg/tonne feed. Feed as sole ration for 14d.

## WITHDRAWAL TIME:

Meat: 7d

## USE:

## Swine dysentery (*Brachyspira* spp), control/prevention

### SPECIES:

---

Swine

### ROA:

---

In the feed

### DOSE:

---

0.386 kg/tonne feed. Feed as sole ration on premises with a history of swine dysentery but where signs of the disease have not yet occurred, or following treatment at 220 mg/kg tiamulin rate to prevent re-infection during risk period of 6-8 weeks.

### WITHDRAWAL TIME:

---

Meat: 2d