



Tiamulin HF 10%

INGREDIENT(S):

tiamulin

USE:

Porcine colonic spirochaetosis (colitis) (*Brachyspira pilosicoli*), treatment
Swine dysentery (*Brachyspira* spp), treatment

SPECIES:

Swine

ROA:

In the feed

DOSE:

2.2 kg/tonne feed. Feed as sole ration for 14d.

WITHDRAWAL TIME:

Meat: 7d

USE:

Porcine proliferative enteritis (PPE, ileitis – *Lawsonia intracellularis*), treatment

SPECIES:

Swine

ROA:

In the feed

DOSE:

1.5 kg/tonne feed. Feed as sole ration for 14d.

WITHDRAWAL TIME:

Meat: 7d

USE:

Pneumonia (enzootic – *Mycoplasma hyopneumoniae*), treatment

SPECIES:

Swine

ROA:

In the feed

DOSE:

1.65 kg/tonne feed. Feed as sole ration for 14d.

WITHDRAWAL TIME:

Meat: 7d

USE:

Swine dysentery (*Brachyspira* spp), control/prevention

SPECIES:

Swine

ROA:

In the feed

DOSE:

0.386 kg/tonne feed. Feed as sole ration on premises with a history of swine dysentery but where signs of the disease have not yet occurred, or following treatment at 220 mg/kg tiamulin rate to prevent re-infection during risk period of 6-8 weeks.

WITHDRAWAL TIME:

Meat: 2d